



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ON A DEFINITION OF ABSTRACT GROUPS*

BY

ELIAKIM HASTINGS MOORE

In a paper entitled *A definition of abstract groups* (Transactions, vol. 3, pp. 485–492, October, 1902), my second definition (l. c., p. 490)

$$(M'') = (1, 2, 3', 3'', 3''_r, 4''_l)$$

involves postulates not mutually independent. I shall prove here (as stated in October, 1904, Transactions, vol. 5, p. 549) that $(3'')$ is redundant, that in the new definition

$$(\bar{M}'') : (1, 2, 3'', 3''_l, 4'')$$

the postulates are mutually independent, and that this mutual independence remains even for the system

$$(\bar{M}_A'') : (1, 2, 3'', 3''_l, 4'', A)$$

defining an abelian group, obtained by adding to those of (M'') the postulate (A) that the multiplication or composition of two elements is commutative.

We have for consideration a set † or class (K) of elements and a multiplication-table or rule of combination $(\circ)$ whereby to every two elements a, b taken in the definite order a, b there corresponds a definite so-called product, in notation $a \circ b$, or, when without confusion, more simply, ab ; this product may or may not be an element of the class. The postulates in question are then the following:

(1) If a and b are elements, then ab is an element of the class.

(A) If a and b are elements such that ab and ba are elements, then $ab = ba$.

(2) If a, b, c are elements such that $ab, bc, (ab)c, a(bc)$ are elements, then $(ab)c = a(bc)$.

(3'') There exists an element a such that $aa = a$.

(3''_l) If a and b are elements and $aa = a$, then $ab = b$.

(3''_r) If a and b are elements and $aa = a$, then $ba = b$.

(4''_l) If a and b are elements and $aa = a$, then there exists an element $b^{(a)}_l$ such that $b^{(a)}_l b = a$.

* Presented to the Society December 30, 1904. Received for publication March 17, 1905.

† In the first definition (M) there was the underlying understanding (l. c., p. 485, footnote ‡) or postulate (0) that the class contain at least one element, and this carried over by implication to (M'') , where however in view of $(3'')$ it was not needed; it is now omitted.

If a is an element such that $aa = a$ we designate those parts of $(3''_l)$, $(3''_r)$, $(4''_l)$ which refer to the element a by $(3''^a_l)$, $(3''^a_r)$, $(4''^a_l)$ respectively. Then we prove that $(3''_r)$ is deducible from $(1, 2, 3'', 3''_l, 4''_l)$ in that we prove that $(3''^a_r)$ is deducible from $(1, 2, 3''^a_l, 4''^a_l)$. This fact is proved by a suitable modification of the method used p. 486, 7° loc. cit. If a and b are elements and $aa = a$, then we designate by b' , b'' elements, which by $(4''^a_l)$ surely exist, such that $b'b = a$, $b''b' = a$ and have in virtue of $(1, 2, 3''^a_l)$ the continued equality,

$$ba = bb'b = abb'b = b''b'bb'b = b''ab'b = b''b'b = ab = b,$$

that is, $ba = b$, which was to be proved. *

The proof that the postulates of $(\bar{M}'_A)$ are independent covers the independence of the postulates of $(\bar{M}'')$ and appears from the following proof systems $(K, \circ)$:

For (1). K = all integers $0, \pm 1, \pm 2, \dots$ except ± 1 . $\circ = +$.

For (2). K = an element a and any class of (at least three) elements x distinct from a . $a \circ a = a$. $a \circ x = x \circ a = x$. $x_1 \circ x_2 = a$ if $x_1 \neq x_2$; $x_1 \circ x_1 = x_3$ where, for every x_1 , x_3 is any x except x_1 .

For (3''). K = all positive integers. $\circ = +$.

For (3'_l). K = a class of (at least two) elements x . $x_1 \circ x_2 = a$, where a is a definite element, the same for all pairs x_1, x_2 .

For (4'_l). K = all positive integers and 0. $\circ = +$.

For (A). $(K, \circ)$ = any non-abelian group.

THE UNIVERSITY OF CHICAGO.

* [A second continued equality

$$ba = aba = b''b'ba = b''aa = b''a = b''b'b = ab = b$$

should be noticed.]